

Physical/Occupational Therapy Prescription

Name: _____ Date of Birth: _____

Diagnosis: Pectoralis Major Tear Code: S29.011

Procedure: Pectoralis Major Repair Surgery Date: _____

Instructions: Pectoralis Major Repair Protocol

Range of motion:

- For the first two weeks post-operatively:
 - o Sling at all times.
 - o No range of motion.
- At two weeks post-operatively:
 - o Begin active-assisted motion, progressing towards active range of motion.
 - o Add stretching at end ranges to regain full rotation, but avoid positions of impingement.
 - o No combined abduction and external rotation until six weeks
 - o Please instruct patient and family in a home exercise program.
 - o Goal passive range of motion by six weeks: 140° elevation
 - o Goal passive range of motion by twelve weeks: 60° external rotation.

Strengthening:

- No strengthening prior to 6 weeks post-operatively, 5 pound weight lifting restriction.
- Then begin strengthening of rotator cuff, scapular stabilizers, and deltoid progressing slowly from isometrics (6 weeks) to bands and light resistive strengthening (9 weeks) to weights and heavy resistive strengthening (12 weeks). Limit to 20 pounds until 12 weeks.
- Once strength is progressing, also add a focus upon scapular mechanics and proprioception.
- At 4.5 months post-operatively, progress towards occupation and sport-specific exercises, planning to return to contact sports at 6 months.

Limitations:

- No range of motion and sling immobilization until two weeks post-operatively.
- No combined abduction and external rotation until six weeks post-operatively.
- No strengthening until six weeks post-operatively and progress slowly with strengthening.
- Three- to five-pound weight lifting restriction until six weeks post-operatively.

Please provide with a home exercise program.

Modalities

Heat before and ice after therapy. Electric stimulation and remaining modalities per therapist.

Frequency: 2-3 times/week Duration: 6 weeks

Signature: _____ Date: _____